

# On Perfect Codes in the Johnson Graph <sup>1</sup>

NATALIA SILBERSTEIN, TUVI ETZION {natalys, etzion}@cs.technion.ac.il

Computer Science Dept., Technion - Israel Institute of Technology, Haifa, ISRAEL

**Abstract.** In this paper we consider the existence of nontrivial perfect codes in the Johnson graph  $J(n, w)$ . We present combinatorial and number theory techniques to provide necessary conditions for existence of such codes and reduce the range of parameters in which 1-perfect and 2-perfect codes may exist.

## 1 Introduction

Codes which attain the sphere packing bound are called perfect codes. The Hamming metric and the Johnson metric are the most important metrics in coding theory on which perfect codes are defined. While for the Hamming space all perfect codes over finite fields are known [1], in the Johnson space it was conjectured by Delsarte in 1970's [2, p. 55] that there are no nontrivial perfect codes. The general nonexistence proof still remains an open problem, although many attempts to solve the problem were made, e.g. [3–10].

The *Johnson space*  $V_w^n$  consists of all  $w$ -subsets of a fixed  $n$ -set  $N = \{1, 2, \dots, n\}$ , for given positive integers  $n$  and  $w$  such that  $0 \leq w \leq n$ . With the Johnson space we associate the *Johnson graph*  $J(n, w)$  with the vertex set  $V_w^n$ , where two  $w$ -subsets are adjacent if and only if their intersection is of size  $w - 1$ . A code  $C$  of such  $w$ -subsets is called an *e-perfect code* in  $J(n, w)$  if the  $e$ -spheres with centers at the codewords of  $C$  form a partition of  $V_w^n$ . In other words,  $C$  is an  $e$ -perfect code if for each element  $v \in V_w^n$  there exists a unique codeword  $c \in C$  such that the distance (in the graph) between  $v$  and  $c$  is at most  $e$ .

A code  $C$  in  $J(n, w)$  can be described as a collection of  $w$ -subsets of  $N$ , but it can be also described as a binary code of length  $n$  and constant weight  $w$ . From a  $w$ -subset  $S$  we construct a *characteristic* binary vector of length  $n$  and weight  $w$  with *ones* in the positions of  $S$  and *zeroes* in the positions of  $N \setminus S$ . The *Johnson distance* between two  $w$ -subsets is half of the number of coordinates in which their characteristic vectors differ. In the sequel we will use a mixed language of sets and binary vectors.

There are three families of trivial perfect codes in  $J(n, w)$ :  $V_w^n$  is 0-perfect; any  $\{v\}$ ,  $v \in V_w^n$ ,  $w \leq n - w$ , is  $w$ -perfect; and if  $n = 2w$ ,  $w$  odd, any pair of disjoint  $w$ -subsets is  $e$ -perfect with  $e = \frac{1}{2}(w - 1)$ .

---

<sup>1</sup>This work was supported in part by the Israel Science Foundation (ISF), Jerusalem, Israel, under Grant 230/08.

In this paper we are interested in three problems concerning perfect codes in  $J(n, w)$ : the existence of 1-perfect codes, the existence of perfect codes in  $J(2w, w)$ , and improving the Roos bound [5] given by:

**Theorem 1.** *If an  $e$ -perfect code in  $J(n, w)$ ,  $n \geq 2w$ , exists, then*

$$n \leq (w-1) \frac{2e+1}{e}.$$

The rest of this paper is organized as follows. In Section 2 we present new divisibility conditions, which are based on the connection between perfect codes in  $J(n, w)$  and block designs. Based on this connection we introduce an improvement of the Roos bound for 1-perfect codes. In Section 3 we examine 2-perfect codes in  $J(2w, w)$  and present necessary conditions for existence of such codes, using Pell equation. Finally in Section 4 we summarize our results.

## 2 One-perfect codes in $J(n, w)$

In this section we consider first the connection between perfect codes in the Johnson graph and block designs.

Let  $t, n, w, \lambda$  be integers with  $n > w \geq t$  and  $\lambda > 0$ , and let  $N$  be an  $n$ -set. A  $t - (n, w, \lambda)$  design is a collection  $C$  of distinct  $w$ -subsets (called *blocks*) of  $N$  with the property that any  $t$ -subset of  $N$  is contained in exactly  $\lambda$  blocks of  $C$ . Clearly,  $C$  is a code in  $J(n, w)$ . On the other hand, the largest  $t$  for which a code  $C$  in  $J(n, w)$  is a  $t$ -design is called the *strength* of the code.

In the sequel we assume w.l.o.g. that  $n = 2w + a$  for some  $a \geq 0$ , since the complement of an  $e$ -perfect code in  $J(n, w)$  is an  $e$ -perfect code in  $J(n, n-w)$  [7].

**Lemma 1.** [9] *If the code  $C$  in  $J(2w + a, w)$  has strength  $\varphi$  then for each  $t$ ,  $0 \leq t \leq \varphi$ , it is a  $t$ -design  $t - (2w + a, w, \lambda_t)$  with*

$$\lambda_t = \frac{\binom{2w+a-t}{w-t}}{\Phi_e(w, a)},$$

where  $\Phi_e(w, a)$  is the size of the sphere with radius  $e$ .

The strength of a possible  $e$ -perfect code  $C$  in  $J(2w + a, w)$  can be used to exclude its existence. We define the polynomial

$$\sigma_e(w, a, t) = \sum_{i=0}^e (-1)^i \binom{t}{i} \sum_{j=0}^{e-i} \binom{w-i}{j} \binom{w+a-t+i}{i+j}. \quad (1)$$

It was proved in [8] that  $C$  is an  $e$ -perfect code in  $J(2w + a, w)$  with strength  $\varphi$  if  $\varphi$  is the smallest positive integer for which  $\sigma_e(w, a, \varphi + 1) = 0$ .

The expression for the strength of an 1-perfect code can be easily calculated from (1). We use this expression to obtain divisibility conditions for such codes and to improve the Roos bound on the length of 1-perfect codes in the Johnson graph.

**Theorem 2.** *Assume there exists an 1-perfect code  $C$  in  $J(2w + a, w)$  with strength  $\varphi = w - d$  for some  $d \geq 0$ . Then  $d > 1$ ,  $d \equiv 0$  or  $1 \pmod{3}$ ,  $w - d \equiv 0, 1, 4$ , or  $9 \pmod{12}$ , and*

$$\frac{\prod_{i=0}^{d-2} (wd - (d + i(d-1)))}{(d-1)!(d-1)^{d-1}d(w-d+1)} \in \mathbb{Z}. \quad (2)$$

*Proof.* Assume that there exists an 1-perfect code  $C$  in  $J(2w + a, w)$ . Therefore, by (1), the strength of  $C$  is  $w - d = \frac{2w+a-1-\sqrt{(a+1)^2+4(w-1)}}{2}$ . Hence it follows that  $d > 1$  and

$$a = \frac{w - d^2 + d - 1}{d - 1}. \quad (3)$$

In [8] it was proved that if there exists an 1-perfect code in  $J(n, w)$  then either  $w \equiv n - w \equiv 1 \pmod{12}$  or  $w \equiv n - w \equiv 7 \pmod{12}$ . In particular,  $w \equiv 1 \pmod{6}$ , 12 divides  $a$ , and hence by (3), 6 divides  $d^2 - d$ . Therefore,  $d \equiv 0$  or  $1 \pmod{3}$ , i.e.,  $d \equiv 0, 1, 3, 4, 6, 7, 9$  or  $10 \pmod{12}$ . Since 12 divides  $a$ , from (3) it follows that  $w \equiv d^2 - d + 1 \pmod{12}$ , i.e.,  $w - d \equiv (d-1)^2 \pmod{12}$ . Therefore,  $w - d \equiv 0, 1, 4$ , or  $9 \pmod{12}$ .

Now, using (3) we write the size of the sphere with radius 1 as follows:  $\Phi_1(w, a) = 1 + w(w+a) = (w+a+d)(w-d+1)$ . By Lemma 1, for  $t = w - d$  we have  $\lambda_{w-d} = \frac{\binom{w+a+d}{w+a}}{(w+a+d)(w-d+1)} = \frac{\binom{w+a+d-1}{d-1}}{d(w-d+1)} \in \mathbb{Z}$ . We simplify the last expression, by using  $w + a + d - 1 = \frac{wd-d}{d-1}$  (implied by (3)), and obtain the divisibility condition (2).  $\square$

By Theorem 1 it follows that if an 1-perfect code exists in  $J(2w + a, w)$ , then  $2w + a \leq 3(w-1)$  and thus  $a \leq w-3$ . We use the divisibility condition (2) in Theorem 2 in order to improve this bound.

**Theorem 3.** *If an 1-perfect code exists in  $J(2w + a, w)$ , then  $a < \frac{w}{11}$ .*

*Proof.* Assume that there exists an 1-perfect code  $C$  with strength  $w - d$  in  $J(2w + a, w)$ . We examine the divisibility condition (2) for several values of  $d$  which are not pruned out by Theorem 2.

- Assume  $d = 3$ . By (2) we have that  $\frac{(3w-3)(3w-5)}{2!2^23(w-2)} = \frac{(w-1)(3w-5)}{8(w-2)} \in \mathbb{Z}$ , which is impossible since  $\text{g.c.d.}(w-1, w-2) = 1$  and  $\text{g.c.d.}(3w-5, w-2) = 1$ . Hence,  $d > 3$ .

- Assume  $d = 4$ . By (2) we have that  $\frac{4(w-1)(4w-7)2(2w-5)}{3!3^34(w-3)} \in \mathbb{Z}$ . Since  $\text{g.c.d.}(w-3, w-1) \in \{1, 2\}$ ,  $\text{g.c.d.}(w-3, 4w-7) \in \{1, 5\}$ , and  $\text{g.c.d.}(w-3, 2w-5) = 1$ , it follows that  $w-3 \leq 2 \cdot 5$ . But by (3) we have  $a = \frac{w-13}{3}$ , and hence  $w > 13$ . Thus,  $d > 4$ .
- Assume  $d = 6$ . By (2) we have that  $\frac{6(w-1)(6w-11)(6w-16)(6w-21)(6w-26)}{5!5^56(w-5)} \in \mathbb{Z}$ . Examining the g.c.d. of  $w-5$  with each factor of the nominator, we obtain that all possible factors of  $w-5$  are from  $\{2, 3, 7, 19\}$ . Since  $w \equiv d^2 - d + 1 \equiv 7 \pmod{12}$ , i.e.,  $w-5 \equiv 2 \pmod{12}$  we obtain that  $w-5 = 2 \cdot 7$ , or  $w-5 = 2 \cdot 19$ , or  $w-5 = 2 \cdot 7 \cdot 19$ . Therefore  $w = 19$ , 43 or 271. By (3) we have that  $a = \frac{w-31}{5}$ , and therefore the only possible value for  $w$  is 271 and  $a = 48$ . But  $\Phi_1(w, a)$  does not divide  $\binom{2w+a-(w-7)}{w+a}$  which contradicts Lemma 1. Thus,  $d > 6$ .

Similarly we obtain contradiction for  $d = 7$  and  $d = 9$  and hence  $d \geq 10$ .

- Assume  $d = 10$ . By (2) we have that  $\frac{10(w-1)(10w-19)(10w-28)(10w-37)(10w-46)(10w-55)(10w-64)(10w-73)(10w-82)}{9! \cdot 9^9 \cdot 10(w-9)} \in \mathbb{Z}$ . Examining the g.c.d. of  $w-9$  with each factor of the nominator, we obtain that all possible factors of  $w-9$  are from  $\{2, 7, 11, 13, 17, 31, 53, 71\}$ . By using that  $w-9 \equiv -2 \pmod{12}$  since  $w \equiv d^2 - d + 1 \equiv 7 \pmod{12}$ , and examining the divisibility condition of Lemma 1 we get the contradiction.

Hence, since  $d \equiv 0, 1 \pmod{3}$ , it follows that  $d \geq 12$ . Thus, by (3) we have

$$a \leq \frac{w - 12^2 + 12 - 1}{11} = \frac{w - 133}{11} < \frac{w}{11}.$$

□

Clearly, as the value of  $d$  is growing, considering the divisibility condition (2) becomes more complicated. But, the same method can be used for further improving the bound of Theorem 3.

### 3 Two-perfect codes in $J(2w, w)$

In this section we show the necessary conditions for the existence of a 2-perfect code in  $J(2w, w)$  using Pell equation and prove that there are no 2-perfect codes in  $J(2w, w)$  for  $w \leq 1.97 \times 10^{7655}$ .

**Theorem 4.** *If a 2-perfect code  $C$  exists in  $J(2w, w)$ , then there is an integer  $m \geq 0$  such that*

$$(c.1) \quad w = \frac{(1+\sqrt{2})^{2m+1} + (1-\sqrt{2})^{2m+1} + 6}{4}, \text{ and}$$

$$(c.2) \quad \gamma \triangleq \sqrt{2}((1+\sqrt{2})^{2m} - (1-\sqrt{2})^{2m}) + 1 \text{ is a square of some integer.}$$

*Proof.* Assume  $C$  is a 2-perfect code in  $J(2w, w)$ . By (1) the strength of such code is

$$\frac{1}{2}(-1 + 2w - \sqrt{8w - 11 \pm 4\sqrt{5 - 6w + 2w^2}}). \quad (4)$$

Hence, there exists an integer  $y$  such that  $y^2 = 5 - 6w + 2w^2$  and thus  $(2w - 3)^2 - 2y^2 = -1$ . Let  $x = 2w - 3$ , and consider the equation

$$x^2 - 2y^2 = -1.$$

This equation is known as Pell equation [11] and it has a family of solutions given by

$$x = \frac{(1 + \sqrt{2})^{2m+1} + (1 - \sqrt{2})^{2m+1}}{2} \quad (5)$$

$$y = \frac{(1 + \sqrt{2})^{2m+1} - (1 - \sqrt{2})^{2m+1}}{2\sqrt{2}} \quad (6)$$

for some integer  $m \geq 0$ . Hence

$$w = \frac{(1 + \sqrt{2})^{2m+1} + (1 - \sqrt{2})^{2m+1} + 6}{4},$$

which completes the proof of (c.1).

Equation (4) implies also that

$$\sqrt{8w - 11 \pm 4\sqrt{5 - 6w + 2w^2}} \in \mathbb{Z}.$$

We distinguish between two cases:

**Case 1:**  $\sqrt{8w - 11 + 4\sqrt{5 - 6w + 2w^2}} \in \mathbb{Z}$ . In this case, there exists an integer  $\alpha$  such that  $\alpha^2 = 8w - 11 + 4\sqrt{5 - 6w + 2w^2} = 8w - 11 + 4y = 4(x + y) + 1$ .

**Case 2:**  $\sqrt{8w - 11 - 4\sqrt{5 - 6w + 2w^2}} \in \mathbb{Z}$ . In this case, there exists an integer  $\beta$  such that  $\beta^2 = 8w - 11 - 4\sqrt{5 - 6w + 2w^2} = 8w - 11 - 4y = 4(x - y) + 1$ .

In other words, at least one of the expressions  $4(x + y) + 1$ ,  $4(x - y) + 1$  should be a square of some integer. By (5) and (6) we obtain:

$$\begin{aligned} \alpha^2 &= 4(x + y) + 1 = \sqrt{2}((1 + \sqrt{2})^{2m+2} - (1 - \sqrt{2})^{2m+2}) + 1, \\ \beta^2 &= 4(x - y) + 1 = \sqrt{2}((1 + \sqrt{2})^{2m} - (1 - \sqrt{2})^{2m}) + 1, \end{aligned}$$

and the theorem is proved.  $\square$

We examine now condition (c.2) of Theorem 4 for  $1 \leq m \leq 10000$ . The only values of  $m$  which satisfy (c.2) are 0, 1, and 2, where  $\gamma = 1, 9$ , and 49, respectively. The corresponding values of  $w$  are 2, 5, and 22, respectively. For  $w = 2$  it can be readily verified that there is no 2-perfect code in  $J(4, 2)$ . It was proved in [8] that if a 2-perfect code exists in  $J(2w, w)$ , then  $w \equiv 2, 26$ , or  $50 \pmod{60}$ . Hence there are no 2-perfect codes for  $w = 5$  and  $w = 22$ . Thus for  $1 \leq w \leq 1.97 \times 10^{7655}$  (considering  $m = 10000$ ), there is no 2-perfect code in  $J(2w, w)$ .

## 4 Conclusion

We have proved the nonexistence of perfect codes in  $J(n, w)$  for large set of new parameters. Further techniques and some nonexistence results can be found in [12].

## References

- [1] F.J. MacWilliams and N.J.A. Sloane, *The Theory of Error-Correcting Codes*, North-Holland, Amsterdam, 1977.
- [2] P. Delsarte, An algebraic approach to association schemes of coding theory, *Philips J. Res.*, **10**, 1–97, 1973.
- [3] E. Bannai, Codes in bi-partite distance-regular graphs, *J. London Math. Soc.*, **2**, 197–202, 1977.
- [4] P. Hammond, On the non-existence of perfect codes and nearly perfect codes, *J. Discrete Math.*, **39**, 105–109, 1982.
- [5] C. Roos, A note on the existence of perfect constant weight codes, *J. Discrete Math.*, **47**, 121–123, 1983.
- [6] W.J. Martin, Completely regular subsets *Ph. D. dissertation, Univ. Waterloo*, 1992.
- [7] T. Etzion, On the nonexistence of perfect codes in the Johnson scheme, *SIAM J. Discrete Math.*, **9**, 201–209, 1996.
- [8] T. Etzion and M. Schwartz, Perfect constant-weight codes, *IEEE Trans. on Inform. Theory*, **50**, 2156–2165, 2004.
- [9] T. Etzion, Configuration distribution and designs of codes in the Johnson scheme, *J. of Combinatorial designs*, **15**, 15–34, 2006.
- [10] D.M. Gordon, Perfect single error-correcting codes in the Johnson scheme, *IEEE Transactions on Information Theory*, **52**, 4670–4672, 2006.
- [11] I. Niven, H.S. Zuckerman, *An Introduction to the Theory of Numbers*, J. Wiley & sons inc., New York, 1972.
- [12] N. Silberstein, Properties of Codes in the Johnson Scheme, *M.Sc. Thesis*, 2007, (in arXiv:1004.4882).